

reserved on STN

AN 2004444159 EMBASE

TI Involvement of inducible costimulator in the exaggerated memory B cell and plasma cell generation in systemic lupus erythematosus.

AU Hutloff A.; Buchner K.; Reiter K.; Baelde H.J.; Odendahl M.; Jacobi A.; Dorner T.; KroczeK R.A.

CS Dr. R.A. KroczeK, Molecular Immunology, Robert Koch Institute, Nordufer 20, D-13353 Berlin, Germany. kroczeK@rki.de

SO Arthritis and Rheumatism, (2004) Vol. 50, No. 10, pp. 3211-3220. .

Refs: 42

ISSN: 0004-3591 CODEN: ARHEAW

CY United States

DT Journal; Article

FS 005 General Pathology and Pathological Anatomy  
026 Immunology, Serology and Transplantation  
028 Urology and Nephrology  
031 Arthritis and Rheumatism

LA English

SL English

ED Entered STN: 12 Nov 2004

Last Updated on STN: 12 Nov 2004

AB Objective. In systemic lupus erythematosus (SLE), the increased generation of memory B cells and plasma cells leads to autoimmune hypergammaglobulinemia and destructive immunoglobulin deposits in the kidneys. We undertook this study to determine the biologic mechanism driving this overactivation of the B cell compartment, which is the central issue in SLE. Methods. We used flow cytometry to analyze expression of the T cell-specific inducible costimulator (ICOS) and its ligand (ICOS-L) on B cells obtained from the peripheral blood of SLE patients. We correlated ICOS-L expression with the differentiation status of the B cells using a large panel of surface antigens. In addition, SLE kidneys were analyzed by immunohistology. Results. We found an increased expression of ICOS on CD4+ as well as CD8+ T cells in SLE. At the same time, we documented a down-regulation of ICOS-L on a high proportion of peripheral blood memory B cells. Based on in vitro experiments, we inferred that this ICOS-L down-regulation on B cells was a signature of recent interaction with ICOS+ T cells in vivo. In the kidneys of SLE patients, we found clusters of B cells and plasma cells in close contact with ICOS+ T cells. Conclusion. Detailed analysis of B cells with down-regulated ICOS-L suggests that ICOS is one of the forces driving the formation of memory B cells and plasma cells in SLE. Furthermore, our identification of plasma cells in areas of T cell-B cell interaction in kidneys suggests that components of a T cell-driven B cell activation process may take place in peripheral tissues in SLE.

=> D His

(FILE 'HOME' ENTERED AT 13:09:56 ON 10 JAN 2007)

FILE 'MEDLINE, CAPLUS, BIOSIS, EMBASE, SCISEARCH' ENTERED AT 13:10:41 ON 10 JAN 2007

L1 E FINNEY HELENE/AU  
27 S E2, E3, E4, E5  
E LAWSON ALASTAIR/AU

L2 56 S E3, E4, E5, E6

L3 64 S L1 OR L2

L4 998 S ICOS (S) ("INDUCIBLE COSTIMULATOR" OR "INDUCIBLE CO-STIMULATOR

L5 1158 S CD134

L6 9 S L4 AND L5

L7 2 S L3 AND L4

L8 9 S L6 OR L7

L9 88 S ("CYTOPLASMIC SIGNALLING DOMAIN" OR "CYTOPLASMIC SIGNALLING S

L10 0 S L4 AND L9

L11 1 S L5 AND L9  
L12 31385 S CD28  
L13 4 S L12 AND L9  
L14 4 S L11 OR L13  
L15 13 S L8 OR L14  
L16 9 DUP REM L15 (4 DUPLICATES REMOVED)

=> Logoff

ALL L# QUERIES AND ANSWER SETS ARE DELETED AT LOGOFF

LOGOFF? (Y)/N/HOLD:Y

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

103.79

104.00

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE

TOTAL

ENTRY

SESSION

CA SUBSCRIBER PRICE

-3.90

-3.90

STN INTERNATIONAL LOGOFF AT 13:24:24 ON 10 JAN 2007